

Table 4A. Cancer Incidence per 100,000 Women, White and Minority, 1992–1998^a

	All Cancers	Colon and Rectum	Lung and Bronchus	Breast	Cervix
Alaska Native ^{b,c}	400.1	76.1	57.8	118.1	7.5
American Indian ^{b,c}	140.1	13.5	12.4	36.2	6.2
Asian and Pacific Islander	252.1	31.0	22.7	78.1	10.3
Black	337.6	44.7	45.7	101.5	11.0
Hispanic	237.7	23.2	18.7	68.5	14.4
White	354.4	36.3	43.6	115.5	8.1

^a Age-adjusted to the 1970 U.S. standard population.

^b Data for Alaska Native women derived from American Indian and Alaska Native in the state of Alaska.

^c Data for American Indian women derived from American Indian and Alaska Native in 11 SEER areas (Alaska excluded).

Source: Surveillance, Epidemiology, and End Results (SEER) Program of the National Cancer Institute (August 2000 submission).

Table 4B. Cancer Incidence per 100,000 Women, White and Minority, 1988–1992^a

	All Cancers	Colon and Rectum	Lung and Bronchus	Breast	Cervix
Alaska Native ^{b,c}	348.0	67.4	50.6	78.9	15.8
American Indian ^c	180.0	15.3	5.4	31.6	9.9
Black	326.0	45.5	44.2	95.4	13.2
Chinese	213.0	33.6	25.3	55.0	7.3
Filipino	224.0	20.9	17.5	73.1	9.6
Japanese	241.0	39.5	15.2	82.3	5.8
Native Hawaiian	321.0	30.5	43.1	105.6	9.3
Hispanic	243.0	24.7	19.5	69.8	16.2
White	346.0	38.3	41.5	111.8	8.7

^a Age-adjusted to the 1970 U.S. standard population.

^b Data for Alaska Native women derived from American Indian and Alaska Native in the state of Alaska.

^c American Indian statistics derived from American Indian and Alaska Native cases in the state of New Mexico only.

Source: Miller et al., 1996.